

NPCA Annual Report 2020 – Provider results

University Hospitals Bristol NHS Foundation Trust

Specialist MDT: North Bristol NHS Trust

Disease Presentation

	Specialist MDT	National
No. of men with disease status determined	1780	49027
Percentage of men diagnosed with metastatic disease	13%	13%

Management

	Specialist MDT	National
No. of men diagnosed with low-risk localised disease	140	3055
Percentage of men with low-risk localised disease receiving radical treatment	6%	5%
No. of men diagnosed with locally advanced disease	679	19663
Percentage of men diagnosed with locally advanced disease receiving radical treatment	69%	71%
No. of men diagnosed with M1 (metastases)	224	6136
Percentage of men diagnosed with M1 (metastases) receiving docetaxel	32%	36%

Patient experience of care

	Specialist MDT	National
No. of men diagnosed 1st April 18 - 30th Sept 18 who had radical treatment (surgery or radiotherapy) and were sent a questionnaire	358	10750
Percentage of men who responded	75%	78%
No. of men who completed CNS question	260	8137
Percentage of men who were 'given the name of a CNS'	90%	87%
No. of men who rated their overall care on a scale of 1-10	255	7878
Percentage of men who rated their overall care as 8 or above	94%	91%


Outcome

	Treatment Centre
No. of men who received radical radiotherapy ()	130
Adjusted percentage of men receiving a procedure of the large bowel and a diagnosis indicating radiation toxicity (GI complication) (%)	8%

Patient-reported outcomes


	Treatment Centre
No. of men diagnosed 1st April 18 - 30th Sept 18 who received external beam radiotherapy and were sent a questionnaire	109
Percentage of men who responded	73%
No. of men who completed sufficient information for a validated EPIC-26 bowel function score to be determined	65
Adjusted EPIC-26 bowel function score (0=worst to 100=best)	90
No. of men who completed sufficient information for a validated EPIC-26 sexual function score to be determined	75
Adjusted EPIC-26 sexual function score (0=worst to 100=best)	15

Adjusted funnel plot for the proportion of patients receiving a procedure of the large bowel and a diagnosis indicating radiation toxicity (gastrointestinal complication) within 2 years of radical radiotherapy by radiotherapy centre in England and Wales.




Trust	Number	Adjusted
University Hospitals Bristol NHS Foundation Trust	130	8.32%

Adjusted funnel plot for the mean sexual function score after radical radiotherapy by radiotherapy centre in England and Wales.



Trust	Number	Adjusted mean score
University Hospitals Bristol NHS Foundation Trust	75	15.09

Adjusted funnel plot for the mean bowel function score after radical radiotherapy by radiotherapy centre in England and Wales.



Trust	Number	Adjusted mean score
University Hospitals Bristol NHS Foundation Trust	65	90.08