

### DIAGNOSIS AND STAGING

# 42,975

men were diagnosed with prostate cancer in England and Wales between 1st April 2016 and 31st March 2017

England      Wales

**80%**      **41%**

of men had a pre-biopsy multiparametric MRI

**12%**      **4%**

of men had a transperineal biopsy


**55%**

of men were 70 years or older


**16%**

of men presented with metastatic disease – no change from 15/16

### TREATMENT ALLOCATION



Fewer men with low-risk, localised disease had radical treatments and were potentially 'over-treated'




Slightly more men with locally-advanced disease did not have radical treatments and were potentially 'under-treated'

### TREATMENT OUTCOMES

**13%**

of men were **readmitted** within 3 months following surgery



Within **2 years of treatment** 1 in 10 men experienced a **severe genitourinary complication after surgery** or a **severe gastrointestinal complication after external beam radiation**

After surgery, men reported their **sexual function** to be **23** and **urinary continence** to be **71** on a scale of 1 to 100

After external beam radiation, men reported their **sexual function** to be **17** and **bowel function** to be **85** on a scale of 1 to 100

### PATIENT EXPERIENCE OF CARE



**90%**

of men said they were **given the 'right amount' of information**



**83%**

of men said they were **'given the name of a clinical nurse specialist'**

**72%**

of men said they were **involved as much as they wanted to be in treatment decision making**

**89%**

of men rated their care as **8/10**